

# Comment on "Minimal conductivity in graphene: Interaction corrections and ultraviolet anomaly" by Mishchenko E. G.

IGOR F. HERBUT<sup>1</sup>, VLADIMIR JURIČIĆ<sup>1</sup>, OSKAR VAFEK<sup>2</sup> and MATTHEW J. CASE<sup>2</sup>

<sup>1</sup> *Department of Physics, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada*

<sup>2</sup> *National High Magnetic Field Laboratory and Department of Physics, Florida State University, Tallahassee, Florida 32306, USA*

PACS 73.23.-b – Electronic transport in mesoscopic systems  
PACS 73.25.+i – Surface conductivity and carrier phenomena

**Abstract.** -

In a recent Letter [1], Mishchenko has presented a calculation of the minimal conductivity of the clean graphene taking into account the  $1/r$  Coulomb interactions to first order in perturbation theory, and questioned the results obtained earlier by us [2]. He performed this calculation in three different ways and obtained three different results unless, as he claimed, a proper short distance cutoff procedure restricting large momentum transfers is implemented.

In this comment we wish to clarify and correct some of the statements made in [1].

First, contrary to the claims in [1] the conductivity calculated using the Kinetic equation is identical to the conductivity calculated from the Kubo formula. When calculating transport properties, it is crucial to ensure that the conservation laws are built into the approximation used for the vertex. For the self-energy calculated at the Hartree-Fock level, the four particle correlators must be calculated by summing the RPA bubble and ladder diagrams (see Fig. 1 in [1]). Such non-perturbative treatment, while strictly speaking uncontrolled, does reproduce the leading order perturbative expansion and does not violate Ward identities associated with the standard conservation laws. In general, however, it does violate RG scaling laws. To calculate the electrical conductivity, it is useful to define (in the reciprocal space)

$$\Lambda_\mu(\mathbf{k}; i\Omega_n) \equiv \frac{1}{\beta} \sum_{i\omega_n} \Pi_\mu(\mathbf{k}, i\omega_n; \mathbf{k}, i\omega_n + i\Omega_n), \quad (1)$$

where  $\Omega_n = 2n\pi/\beta$  is a bosonic Matsubara frequency and  $\beta = 1/T$ , and the vertex matrix in the real space (and imaginary time) is

$$\Pi_\mu(x - y, y - x') = -\langle T_\tau \psi(x) \psi^\dagger(x') J_\mu(y) \rangle. \quad (2)$$

Setting  $e = v_F = \hbar = 1$ , the Dirac particle electrical current operator is  $J_\mu(y) = \psi^\dagger(y) \sigma_\mu \psi(y)$ . Within the previously mentioned approximation we have

$$\begin{aligned} \Lambda_\mu(\mathbf{k}; i\Omega_n) = & -\frac{1}{\beta} \sum_{i\omega_n} G_\mathbf{k}(i\omega_n) \sigma_\mu G_\mathbf{k}(i\omega_n + i\Omega_n) \\ & - \int \frac{d^2\mathbf{p}}{(2\pi)^2} V_{\mathbf{k}-\mathbf{p}} \frac{1}{\beta} \sum_{i\omega_n} G_\mathbf{k}(i\omega_n) \Lambda_\mu(\mathbf{p}; i\Omega_n) G_\mathbf{k}(i\omega_n + i\Omega_n). \end{aligned} \quad (3)$$

The above corresponds to 4 coupled integral equations. A rather straightforward calculation shows that the vertex matrix function  $\Lambda_\mu(\mathbf{k}, i\Omega_n)$  can be written in the following form

$$\Lambda_\mu(\mathbf{k}; i\Omega) = \left( \delta_{\mu\nu} - \hat{k}_\mu \hat{k}_\nu \right) \sigma_\nu f_\mathbf{k}(i\Omega) + \epsilon_{\mu\nu} \hat{k}_\nu \sigma_3 g_\mathbf{k}(i\Omega). \quad (4)$$

Here,  $\mu$  runs from 1 to 2 and the sum over repeated indices is understood;  $\epsilon_{\mu\nu}$  is the completely antisymmetric rank 2 tensor and  $\hat{k}$  is a unit length wavevector. This decomposition allows us to reduce the set of 4 coupled integral equations to a set of 2 coupled integral equations for  $f$  and  $g$  with a non-separable vertex. At  $T = 0$ , this set of equations reads

$$\begin{aligned} \begin{bmatrix} 2\epsilon_\mathbf{k} & -\Omega \\ \Omega & 2\epsilon_\mathbf{k} \end{bmatrix} \begin{bmatrix} f_\mathbf{k}(i\Omega) \\ g_\mathbf{k}(i\Omega) \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \\ + \int \frac{d^2\mathbf{p}}{(2\pi)^2} \hat{\mathbf{k}} \cdot \hat{\mathbf{p}} V_{\mathbf{k}-\mathbf{p}} \begin{bmatrix} \hat{\mathbf{k}} \cdot \hat{\mathbf{p}} f_\mathbf{p}(i\Omega) \\ g_\mathbf{p}(i\Omega) \end{bmatrix} \end{aligned} \quad (5)$$

where the Hartree-Fock energy is given by

$$\epsilon_\mathbf{k} = k + \frac{1}{2} \int \frac{d^2\mathbf{p}}{(2\pi)^2} \hat{\mathbf{k}} \cdot \hat{\mathbf{p}} V_{\mathbf{k}-\mathbf{p}}, \quad (6)$$

and (restoring the physical units) the Coulomb potential is

$$V_{\mathbf{k}} = \frac{2\pi}{\epsilon} \frac{e^2}{\hbar v_F} \frac{1}{|\mathbf{k}|}. \quad (7)$$

Eqs. (5) constitutes the leading approximation to the quantum transport theory for massless Dirac fermions in the collisionless limit. The electrical conductivity<sup>1</sup> for  $N$  two components Dirac flavors, in units of  $Ne^2/h$ , can be calculated using

$$\sigma = -2\pi \lim_{\Omega \rightarrow 0} \frac{1}{\Omega} \int \frac{d^2\mathbf{k}}{(2\pi)^2} \Im m f(\mathbf{k}; \Omega + i0^+). \quad (8)$$

The Figure shows the numerical result found perturbatively for small  $\alpha = e^2/(\epsilon\hbar v_F)$  which we compare with the analytic result found to first order in  $\alpha$ .

Second, and more importantly, while the conductivity calculated using Kubo formula would superficially indeed seem to depend on the cutoff procedure, further analysis shows, of course, that it does not. This point can be amply illustrated by using *two* different cutoffs: one hard cutoff  $\Lambda$  which restricts the fermion modes to the vicinity of the Dirac point, and second  $M$  which restricts the momentum transfer, i.e.,

$$V_{\mathbf{k}} \rightarrow \frac{2\pi}{\epsilon} \frac{e^2}{\hbar v_F} \left( \frac{1}{|\mathbf{k}|} - \frac{1}{\sqrt{\mathbf{k}^2 + M^2}} \right). \quad (9)$$

In the high energy parlance the second way of cutting off the theory corresponds to the Pauli-Villars regularization. The result obtained using this procedure is the dc conductivity

$$\sigma = N \frac{e^2}{h} \left[ \frac{\pi}{8} + \alpha \frac{\pi}{16} \left( \frac{25}{6} - \pi - \frac{1}{\sqrt{1 + \frac{M^2}{\Lambda^2}}} \right) \right] \quad (10)$$

where  $\alpha = e^2/\hbar v_F$  and  $N = 4$  is the number of fermion flavors in graphene. Naively, this would seem to lead to a one parameter family of results that interpolate between our original result (recovered when  $M/\Lambda \rightarrow \infty$ ) and the one advocated by Mishchenko (when  $M/\Lambda \rightarrow 0$ ). The root of this apparent ambiguity may be easily understood by considering the limit  $M \gg \Lambda$ , for example. One finds,

$$\sigma \approx \frac{e^2}{h} \left[ \frac{\pi}{8} + \alpha \frac{\pi}{16} \left( \frac{25}{6} - \pi - \frac{\Lambda}{M} \right) \right].$$

The last term proportional to  $\alpha\Lambda/M$  thus corresponds to the perturbatively *irrelevant* short range part of the interaction. Indeed, if one would compute the correction

<sup>1</sup>The conductivity can be calculated without analytical continuation by subtracting the zero frequency component of  $f$  before taking the limit  $\Omega \rightarrow 0$ :

$$\sigma = -2\pi \lim_{\Omega \rightarrow 0} \frac{1}{\Omega} \int \frac{d^2\mathbf{k}}{(2\pi)^2} \{f(\mathbf{k}; i\Omega) - f(\mathbf{k}; 0)\}.$$




Fig. 1: (Upper panel) This is the result of the numerical solution of the integral equation (5), integrated over  $dkk$  for different values of  $e^2$ , ( $v_F = 1, \epsilon = 1$ ), and (Euclidean)  $\Omega$ . The analytical result for the  $e^2 = 0$ ,  $(\frac{1}{2} - \frac{\Omega}{4} \tan^{-1} \frac{2}{\Omega})$  is also plotted with a solid line. The agreement at  $e^2 = 0$  is to the 6th significant digit. The  $\Omega$ -slope of these lines is found at different values of  $\Omega$  and plotted vs.  $e^2$  (not shown). (Lower panel) The  $\alpha$ -slope is then plotted for different values of  $\Omega$ . The red dot is the value found analytically. The deviation at small frequency is likely a result of numerical errors.

to the Gaussian conductivity in the same system of Dirac fermions interacting via short-range interactions only, the result would be

$$\sigma'(\Omega) = \frac{1}{\Omega} [a\lambda\Lambda^2 + b\lambda\Lambda\Omega + c\lambda\Omega^2 + O(\Omega^3, \lambda^2)] \frac{e^2}{h}$$

where  $a$ ,  $b$ , and  $c$  are numerical constants.  $\lambda$  is the dimensionfull short range interaction coupling constant with the dimension one, that can be identified with  $\sim \alpha/M$  in the preceding discussion. The first *two* terms explicitly contain the cutoff, and thus clearly represent artifacts of the calculation that violates gauge invariance. *Both* of these

terms must be dropped, so that the final result becomes

$$\sigma'(\Omega) = [c\lambda\Omega + O(\Omega^2, \lambda^2)] \frac{e^2}{h}.$$

This then agrees with the irrelevance of the short range interaction by power counting. The same result is obtained using the dimensional regularization of Veltman and t'Hooft, which was designed precisely to automatically discard the spurious terms of the type of the above.

The very last term in Eq. (9) therefore is the consequence of the incorrect inclusion of the irrelevant short range interaction into the leading correction to the Gaussian result. Such a leading correction can come only from the single marginally irrelevant coupling in the theory, which is the Coulomb interaction. Short range interaction provides only the further, sub-leading corrections, proportional to the frequency. Kubo formula this way leads to the unique result obtained previously by us [2].

The last point raised by Ref. [1] is that the conductivity calculated using density-density correlation function followed by the continuity equation is divergence free and coincides with the Pauli-Villars result. While there is indeed no logarithmic divergence in vertex correction diagram, the self energy is logarithmically divergent, as it may be readily seen from Eq. (12) in Ref. [1], and must be cut off. We admit that the agreement between these two procedures is puzzling. Furthermore, the Coulomb correction to the dc conductivity calculated using the density-density correlator is not unique, i.e., it depends on the procedure implemented to regularize integrals. Namely, the self-energy diagram when calculated within the dimensional regularization scheme, together with the vertex diagram, yields the correction to the conductivity  $\sigma_{\text{pol}}^{\text{dim-reg}} = \sigma_0 \alpha(11 - 3\pi)/6$ , *different* than one obtained in [1] using sharp momentum cutoff. Here  $\sigma_0 = \pi/2$  is the noninteracting (Gaussian) conductivity in units  $e^2/h$ . The origin of this non-uniqueness is unclear at the moment, but we suspect that it may be the non-gauge invariant contribution to the conductivity which is not properly treated within the density polarization approach. On the other hand, as we have already discussed, the Kubo formula yields a unique, regularization independent Coulomb correction to the dc conductivity found by us in Ref. [2].

## REFERENCES

- [1] Mishchenko E. G., Europhys. Lett. **83**, (2008) 17005.
- [2] Herbut I. F., Juričić V. and Vafek O., Phys. Rev. Lett. **100** (2008) 046403.